

Beat: Health

FOOD ODORS ACTIVATE IMPULSE AREA OF THE BRAIN IN OBESE CHILDREN

OBSESSIVE-COMPULSIVE DISORDER

PARIS -CHICAGO, 25.11.2015, 13:19 Time

USPA NEWS - The area of the brain associated with impulsivity and the development of obsessive-compulsive disorder is activated in obese children when introduced to food smells, according to a study being presented next week at the annual meeting of the Radiological Society of North America (RSNA).

The area of the brain associated with impulsivity and the development of obsessive-compulsive disorder is activated in obese children when introduced to food smells, according to a study being presented next week at the annual meeting of the Radiological Society of North America (RSNA).

In the United States, nearly 12.7 million children are obese, according to the Centers for Disease Control and Prevention (CDC).

These children are at a higher risk to develop high blood pressure, type 2 diabetes, and breathing and joint problems, among many other health issues. They are also more likely to become obese adults.

The researchers studied 30 children between the ages of 6 and 10 years old. Half of the children had a normal body mass index (BMI) between 19 and 24, and the other half exhibited a BMI over 30, which is classified as obese. Each child was presented with three odor samples: chocolate, onion and a neutral odor of diluted acetone.

As the participants smelled the samples, two MRI techniques, functional MRI (fMRI) and functional connectivity MRI (fcMRI), were used to measure brain activity.

An evaluation of the fMRI results showed that in the obese children, the food odors triggered activation in the areas of the brain associated with impulse and the development of obsessive-compulsive disorder, while the areas of the brain associated with impulse control exhibited no activity. However, in the children with a normal BMI, the areas of the brain associated with pleasure regulation, organization and planning, as well as regions governing emotional processing or memory function, became more active.

In addition, the fcMRI results showed that when the normal-weight children smelled the onion, there was a connection between the gustatory cortex, which processes taste, and the area of the brain linked to reward anticipation. This connection did not occur in the obese children.

The chocolate smell elicited significant brain connections in obese children, compared to the normal-weight children.

Co-authors on this study were Silva Hidalgo-Tobon, Ph.D., Benito De Celis IV, Eduardo Barragan, Eduardo Castro, M.D., Samuel Flores, M.D., Porfirio Ibanez and Manuel Obregon.

RSNA is an association of more than 54,000 radiologists, radiation oncologists, medical physicists and related scientists, promoting excellence in patient care and health care delivery through education, research and technologic innovation.

Copies of RSNA 2015 news releases and electronic images will be available online at RSNA.org/press15 beginning Monday, Nov. 30.

Source : Radiological Society of North America (RSNA)

Ruby BIRD

<http://www.portfolio.uspa24.com/>

Yasmina BEDDOU

<http://www.yasmina-beddou.uspa24.com/>

Article online:

<https://www.uspa24.com/bericht-6536/food-odors-activate-impulse-area-of-the-brain-in-obese-children.html>

Editorial office and responsibility:

V.i.S.d.P. & Sect. 6 MDSStV (German Interstate Media Services Agreement): Ruby BIRD & Yasmina BEDDOU (Journalists/Directors)

Exemption from liability:

The publisher shall assume no liability for the accuracy or completeness of the published report and is merely providing space for the submission of and access to third-party content. Liability for the content of a report lies solely with the author of such report. Ruby BIRD & Yasmina BEDDOU (Journalists/Directors)

Editorial program service of General News Agency:

United Press Association, Inc.
3651 Lindell Road, Suite D168
Las Vegas, NV 89103, USA
(702) 943.0321 Local
(702) 943.0233 Facsimile
info@unitedpressassociation.org
info@gna24.com
www.gna24.com